

Protecting and improving the nation's health

Walsall

Unitary Authority



This profile was produced on 2 June 2015

Health Profile 2015

Health in summary

The health of people in Walsall is generally worse than the England average. Deprivation is higher than average and about 27.9% (15,500) children live in poverty. Life expectancy for both men and women is lower than the England average.

Living longer

Life expectancy is 10.1 years lower for men and 7.5 years lower for women in the most deprived areas of Walsall than in the least deprived areas.

Child health


In Year 6, 24.4% (772) of children are classified as obese, worse than the average for England. The rate of alcohol-specific hospital stays among those under 18 was 48.8*. This represents 32 stays per year. Levels of teenage pregnancy, GCSE attainment, breastfeeding and smoking at time of delivery are worse than the England average.

Adult health

In 2012, 30.4% of adults are classified as obese, worse than the average for England. The rate of alcohol related harm hospital stays was 730*, worse than the average for England. This represents 1,864 stays per year. The rate of self-harm hospital stays was 196.0*. This represents 537 stays per year. The rate of smoking related deaths was 320*, worse than the average for England. This represents 463 deaths per year. Estimated levels of adult excess weight and physical activity are worse than the England average. Rates of sexually transmitted infections and TB are worse than average. The rate of people killed and seriously injured on roads is better than average.

Local priorities

Priorities in Walsall include reducing infant mortality, promoting healthy weight, tackling alcohol, and tackling health inequalities particularly in men. For more information see 'health and social care' at <u>cms.walsall.gov.uk</u>



Contains OS data @ Crown copyright and database rights 2015 @ OpenStreetMap contributors ODbL

Population: 272,000

Mid-2013 population estimate. Source: Office for National Statistics.

This profile gives a picture of people's health in Walsall. It is designed to help local government and health services understand their community's needs, so that they can work together to improve people's health and reduce health inequalities.


Visit <u>www.healthprofiles.info</u> for more profiles, more information and interactive maps and tools.

Follow <u>@PHE_uk</u> on Twitter

* rate per 100,000 population

Deprivation: a national view


The map shows differences in deprivation in this area based on national comparisons, using quintiles (fifths) of the Index of Multiple Deprivation 2010, shown by lower super output area. The darkest coloured areas are some of the most deprived neighbourhoods in England. This chart shows the percentage of the population who live in areas at each level of deprivation.




Life expectancy: inequalities in this local authority

The charts below show life expectancy for men and women in this local authority for 2011-2013. Each chart is divided into deciles (tenths) by deprivation, from the most deprived decile on the left of the chart to the least deprived decile on the right. The steepness of the slope represents the inequality in life expectancy that is related to deprivation in this local area. If there were no inequality in life expectancy as a result of deprivation, the line would be horizontal.









Health inequalities: changes over time

These charts provide a comparison of the changes in early death rates (in people under 75) between this area and all of England. Early deaths from all causes also show the differences between the most and least deprived quintile in this area. (Data points are the midpoints of 3 year averages of annual rates, for example 2005 represents the period 2004 to 2006).



Health inequalities: ethnicity



Percentage of hospital admissions that were emergencies, by ethnic group, 2013

Health summary for Walsall

The chart below shows how the health of people in this area compares with the rest of England. This area's result for each indicator is shown as a circle. The average rate for England is shown by the black line, which is always at the centre of the chart. The range of results for all local areas in England is shown as a grey bar. A red circle means that this area is significantly worse than England for that indicator; however, a green circle may still indicate an important public health problem.

	cantly worse that			England	Regional average [^]		England Average		England	
		ent from England average			Worst		25th		75th	Best
Significantly better than England average Domain Indicator			Local No Per Year	Local value	Eng value	Eng worst	Percentile	England Range	Percentile	Eng best
Our communities	1 Deprivation		124,940	45.9	20.4	83.8		• •		0.0
	2 Children in p	poverty (under 16s)	15,490	27.9	19.2	37.9				5.8
	3 Statutory ho	omelessness	332	3.1	2.3	12.5				0.0
	4 GCSE achie	eved (5A*-C inc. Eng & Maths)†	1,534	48.5	56.8	35.4				79.9
	5 Violent crim	e (violence offences)	2,902	10.7	11.1	27.8				2.8
	6 Long term u	inemployment	2,363	14.1	7.1	23.5				0.9
Children's and young people's health	7 Smoking sta	atus at time of delivery	487	13.7	12.0	27.5				1.9
	8 Breastfeedir	ng initiation	2,291	63.4	73.9					
	9 Obese child	lren (Year 6)	772	24.4	19.1	27.1		•		9.4
	10 Alcohol-spe	cific hospital stays (under 18)†	31.7	48.8	40.1	105.8		● 🔶		11.2
	11 Under 18 cc	onceptions	192	36.8	24.3	44.0	•			7.6
Adults' health and lifestyle	12 Smoking pre	evalence	n/a	20.5	18.4	30.0				9.0
	13 Percentage	of physically active adults	244	50.7	56.0	43.5		• •		69.7
	14 Obese adult	ts	n/a	30.4	23.0	35.2	•			11.2
	15 Excess weig	ght in adults	463	68.9	63.8	75.9				45.9
Disease and poor health	16 Incidence of	f malignant melanoma†	26.3	11.7	18.4	38.0			•	4.8
	17 Hospital sta	ys for self-harm	537	196.0	203.2	682.7		$\langle \phi \rangle$		60.9
	18 Hospital sta	ys for alcohol related harm†	1,864	730	645	1231				366
	19 Prevalence	of opiate and/or crack use	2,186	12.8	8.4	25.0				1.4
	20 Recorded di	iabetes	18,579	8.6	6.2	9.0	•			3.4
	21 Incidence of	f TB†	56.7	20.9	14.8	113.7				0.0
	22 New STI (ex	xc Chlamydia aged under 25)	1,753	1026	832	3269				172
	23 Hip fracture	s in people aged 65 and over	283	550	580	838				354
Life expectancy and causes of death	24 Excess wint	ter deaths (three year)	169.3	22.3	17.4	34.3		•		3.9
	25 Life expecta	ancy at birth (Male)	n/a	78.1	79.4	74.3				83.0
	26 Life expecta	ancy at birth (Female)	n/a	82.8	83.1	80.0				86.4
	27 Infant morta	ılity	26	6.8	4.0	7.6	•	•		1.1
	28 Smoking rel	lated deaths	463	319.6	288.7	471.6		•		167.4
	29 Suicide rate	1	17	6.5	8.8					
	30 Under 75 m	ortality rate: cardiovascular	209	95.1	78.2	137.0				37.1
	31 Under 75 m	ortality rate: cancer	355	161.5	144.4	202.9				104.0

Indicator notes

1 % people in this area living in 20% most deprived areas in England, 2013 2 % children (under 16) in families receiving means-tested benefits & low income, 2012 3 Crude rate per 1,000 households, 2013/14 4 % key stage 4, 2013/14 5 Recorded violence against the person crimes, crude rate per 1,000 population, 2013/14 6 Crude rate per 1,000 population aged 16-64, 2014 7 % of women who smoke at time of delivery, 2013/14 8 % of all mothers who breastfeed their babies in the first 48hrs after delivery, 2013/14 9 % school children in Year 6 (age 10-11), 2013/14 10 Persons under 18 admitted to hospital due to alcohol-specific conditions, crude rate per 100,000 population, 2011/12 to 2013/14 (pooled) 11 Under-18 conception rate per 1,000 females aged 15-17 (crude rate) 2013 12 % adults aged 18 and over who smoke, 2013 13 % adults achieving at least 150 mins physical activity per week, 2013 14 % adults classified as obese, Active People Survey 2012 15 % adults classified as overweight or obese, Active People Survey 2012 16 Directly age standardised rate per 100,000 population, aged under 75, 2010-12 17 Directly age standardised rate per 100,000 population, 2013/14 18 The number of admissions involving an alcohol-related primary diagnosis or an alcohol-related external cause, directly age standardised rate per 100,000 population, 2013/14 19 Estimated users of opiate and/or crack cocaine aged 15-64, crude rate per 1,000 population, 2011/12 20 % people on GP registers with a recorded diagnosis of diabetes 2013/14 21 Crude rate per 100,000 population, 2013 23 Directly age and sex standardised rate of emergency admissions, per 100,000 population aged 65 and over, 2013/14 24 Ratio of excess winter deaths (observed winter deaths minus expected deaths based on non-winter deaths) to average non-winter deaths 01.08.10-31.07.13 25, 26 At birth, 2011-13 27 Rate per 1,000 live births, 2011-13 28 Directly age standardised rate per 100,000 population aged under 75, 2011-13 30 Directly age standardised rate per 100,000 population aged

† Indicator has had methodological changes so is not directly comparable with previously released values.

^ "Regional" refers to the former government regions.

More information is available at www.healthprofiles.info and https://fingertips.phe.org.uk/profile/health-profiles.

Please send any enquiries to healthprofiles@phe.gov.uk

You may re-use this information (not including logos) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence, visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3/

4 www.healthprofiles.info